

REVIEW

Gene therapy progress and prospects cancer: oncolytic viruses

T-C Liu¹ and D Kirn^{1,2}

¹Department of Clinical Research, Jennerex Biotherapeutics Inc., San Francisco, CA, USA and ²Department of Pharmacology, School of Medicine, University of Oxford, Oxford, UK

The past 2 years have seen several major advances in oncolytic virotherapy. Studies on the interaction between viruses, immune responses and tumor microenvironment have provided important insight, while clinical trials have shown promise. This review summarizes key findings in this

field over the past 2 years, and provides directions for future success.

Gene Therapy (2008) 15, 877–884; doi:10.1038/gt.2008.72; published online 17 April 2008

Keywords: cancer; oncolytic; virus; replication

In brief

Progress

- Suppression of innate immune response enhances efficacy
- Carrier cell strategy avoids immune attack
- Targeting tumor microenvironment enhances viral spread and efficacy
- Oncolytic viruses kill cancer stem cells
- Genetic engineering of oncolytic viruses complements chemo- and molecular-targeted therapies
- Genetic engineering of oncolytic viruses targets cancer signaling pathways
- Novel oncolytic virus species are being explored
- A large number of clinical trials have been carried out

Prospects

- Logical design of the next generation of oncolytic viruses may take this strategy to the next level
- *Ex vivo* studies may predict responses
- The demand for target validation is increasing
- New imaging endpoints in clinical trials.

Introduction

The use of live viruses for the treatment of cancer dates back to a century.¹ Cancer-selective oncolytic viruses replicate preferentially in cancer cells and as a result, destroy those cells at the end of replication cycles; normal cells are spared and hence toxicity is limited. Of note, oncolytic viruses can kill apoptosis-resistant tumor cells, and hence do not have cross-resistance with existing therapies. Engineered oncolytic viruses have been developed over the past 15 years and have various mechanisms-of-actions (MOA; Table 1): (1) inherently tumor-selective virus species (for example, RNA viruses, poxviruses); (2) viral gene-deleted mutants—critical viral gene expendable for growth in tumor cells, but not in

normal cells, were deleted (for example, adenovirus *dl1520/Onyx-015*, herpes simplex virus (HSV) G207); (3) promoter engineered mutants—viral replication was engineered to be dependent on inserted tumor-specific promoters, and as a result, the replication of the virus is restricted to tumor cells that are able to activate the promoters (for example, prostate specific antigen-regulated adenovirus CG7870, telomerase-regulated adenoviruses and HSVs); (4) pseudotyped viruses—normal viral tropism is ablated, and viruses are engineered to attach/bind to specific surface receptors that are expressed exclusively/preferentially on tumor cells (for example, adenovirus Delta-24RGD).

Over the past decade, several oncolytic viruses have been tested in humans, and although the safety results are encouraging, efficacy as single agents was limited.¹ Possible hurdles include attenuation of the virus caused by genetic engineering of the virus that renders cancer selectivity, host immune responses and lack of understanding of tumor microenvironment. However, H101, an oncolytic adenovirus similar to Onyx-015 (E1B-55K/

Correspondence: Dr D Kirn, Jennerex Biotherapeutics Inc., One Market Street, Spear Tower, Suite 2260, San Francisco, CA 94105, USA.

E-mail: dkirn@jennerex.com

Received 14 January 2008; revised 7 March 2008; accepted 10 March 2008; published online 17 April 2008

Table 1 Cancer-selectivity mechanisms of oncolytic viruses

Approach to selectivity	Agent(s) example and genetic alterations within virus	Genetic target(s) in tumors
Inherently tumor-selective species	NDV (none) Reovirus (none) VSV (none)	IFN resistance Ras pathway IFN resistance
Deletion of viral gene that is necessary for replication in normal cells, but expendable in tumor cells	G207 (ICP6-/γ34.5-deleted HSV-1) Onyx-015 (E1B-55K-/E3B-deleted Ad) Delta-24 (E1A-CR2-deleted Ad) JX-594 (TK-deleted VV)	Proliferation, loss of neurovirulence Loss of p53 pathway, late mRNA transport Loss of G ₁ -S checkpoint control; loss of pRB function Proliferation
Tumor-/tissue-specific promoter engineering to limit viral gene expression	CG7870 (E1A under rat probasin promoter, E1B under PSA promoter/enhancer Ad) bM24-TE (Wnt/β-catenin-promoter/enhancer-driven HSV-1)	Prostate cancer Wnt/β-catenin-overexpressing tumors (colorectal, hepatoblastoma and so on)
Pseudotyped viruses	CAR/integrin-binding deleted Ad, replaced with tumor-targeting ligand	Tumor-specific receptor

Abbreviations: CAR, coxsackievirus-adenovirus receptor; IFN, interferon; NDV, Newcastle disease virus; VSV, vesicular stomatitis virus.

E3B-deleted), was recently approved by the Chinese government to be used in conjunction with radiation therapy for the treatment of head and neck cancers. This is the first oncolytic virus product approved by a governmental agency for human use. To overcome the obstacles toward efficacious virotherapeutics, several major advances have been made to improve the selectivity and efficacy of oncolytic viruses. This review summarizes recent major advances over the past 2 years from over 400 publications and selected unpublished work.

Progress

Suppression of innate immune response enhances efficacy

Virus-immune system interactions have been extensively studied in the context of virotherapy. Innate immune responses to the virus are a major hurdle for long-term gene expression and oncolytic potency. The use of immunomodulatory agents in combination with oncolytic viruses was first reported in the 1970s. Recently, several groups have shown, using different viruses and animal models, that administration of cyclophosphamide, known to inhibit innate immune responses, can significantly enhance viral spread, transgene expression and antitumoral efficacy. These studies demonstrated that cyclophosphamide is able to inhibit neutralizing antibody induction, macrophages, regulatory T cells (Tregs) induction and intratumoral interferon (IFN)-γ production.²⁻⁸ It remains to be determined, however, how much this approach will benefit cancer patients who often have already various degrees of 'pre-existing' immunosuppression due to disease and chemotherapy.

In addition, Haralambieva *et al.*⁹ showed that measles virus-induced gene expression and intratumoral virus spread is inhibited by IFN, which is triggered by virus infection of tumor cells. Interestingly, currently available oncolytic measles viruses are derived from Edmonston tag (Edmtag) strain, which has lost most of the IFN-antagonizing activities. Edmtag-based measles virus engineered to express the measles phosphoprotein (P) gene products (P/V/C proteins) from wild-type measles virus, known to antagonize IFN induction and response, exhibited reduced IFN sensitivity and a reduced IFN induction in lymphoma, myeloma and activated peripheral blood mononuclear cells (PBMC).⁹ Measles virus encoding the P gene products also showed significantly enhanced systemic efficacy in a myeloma xenograft model. This study highlights the importance of innate antiviral responses of tumor cells that need to be considered when designing oncolytic viruses.

Importantly, antiviral immunity does not necessarily reduce the efficacy of virus. A recent report by Zhu *et al.*¹⁰ demonstrated that in mice pre-immunized with HSV, subsequent intratumoral administration of oncolytic HSV showed enhanced efficacy compared to HSV-naive mice. The peripheral blood mononuclear cells from the HSV-seropositive mice also exhibited greater *in vitro* cytotoxicity to tumor cells than PBMC from HSV-naive mice, which correlated with an enhanced IFN-γ induction in PBMC from HSV-seropositive mice.¹⁰ This is an important finding that needs to be explored with systemic HSV administration and also with other oncolytic virus species.

Consistent with our previous findings, Diaz *et al.*¹¹ have shown that host CD8 and NK cells are critical for the efficacy of oncolytic virotherapy. Importantly, using an antibody against Treg (PC-61) they showed that depletion of Treg cells inhibited antitumoral efficacy in

the context of virotherapy, as it overcame the suppression of antiviral immune responses. Furthermore, levels of activated T cells can be significantly increased by enhanced expression of tumor antigens. This can be achieved either through adoptive T-cell transfer therapy, or incorporation of tumor antigen into the oncolytic virus.¹¹

Carrier cell strategy avoids immune attack

In addition to blocking the host immune response, one can take advantage of the immune system to boost antitumor responses. Adenovirus and HSV mutants engineered to enhance the expression of class I major histocompatibility complex (MHC) have been shown to enhance enhanced antitumoral immune responses and efficacy in animal models. However, one major challenge for virotherapy is the inefficient uptake of viruses into tumor cells after systemic administration due to systemic antiviral immune response (for example, neutralizing antibodies and complement). Thorne *et al.*¹² described a novel approach to tackle this issue. Cytokine-induced killer (CIK) cells are known to 'home' to and destroy tumors. After isolating the CIK cells from mice, these cells were infected with oncolytic vaccinia viruses and readministered into tumor-bearing animals. The virus replicated in the CIK cells while these traveled to the tumors. As a result, substantially larger amounts of oncolytic viruses were delivered to the tumor, and both the CIK cells and oncolytic viruses were synergistic in tumor killing.¹² It remains to be seen, however, whether CIK cells home to tumors in humans within a reasonable time frame. In addition, this approach requires harvesting of cells from individual patients, *ex vivo* culturing and redelivery to the patients, and therefore requires a substantial amount of laboratory work. Nonetheless, this strategy holds promise.

Subsequently, the 'carrier cell' approach has been tested by several groups.¹³ Using measles viruses, Ong *et al.*¹⁴ showed that virus-infected T cells can protect from low, but not high, concentrations of antimeasles immune serum. However, even in measles-naïve mice, only 1–2% of the virus-infected T cells trafficked to the tumor site after systemic delivery. Power *et al.*¹⁵ tested oncolytic vesicular stomatitis virus (VSV) vectors with different carrier cell types, ranging from leukemia cell lines (which led to systemic delivery of the virus) to cells derived from solid tumors (which accumulated primarily in the lungs). Using dual-enzyme *in vivo* luminescence imaging, it was shown that whereas the carrier cells were retained in the body for no longer than 1 day, oncolytic VSV continued to replicate and was able to eradicate pre-established tumors.¹⁵ Other carrier cells tested include endothelial cells and mesenchymal stem cells.^{16,17} Vile *et al.* have recently shown that T lymphocytes can be used to harbor oncolytic VSV and release the virus at the tumor site.¹⁸ VSV-loaded lymphocytes were also able to purge spleen and lymph nodes of metastatic cells, which in turn primed antitumoral immunity. Furthermore, adoptive transfer of VSV-loaded lymphocytes reduced metastases.¹⁸ This strategy might have great potential for many tumor types.

While the 'Trojan Horse' approach avoids the negative impacts of neutralizing antibodies on virotherapy, there are several critical issues that need to be further investigated before this approach can be taken into humans. First of all, the most appropriate carrier cell

types need to be carefully determined. Tumor-homing cells (for example, CIK) and systemic disseminating cells (for example, leukemic cells) have both advantages and disadvantages. Importantly, tumorigenicity is a major concern when cancer cells are used as carriers. Finally, although viruses are able to replicate, the newly generated viruses (that is, 'second wave') still face contact with neutralizing antibodies, which will likely restrict significant virus spread. Multiple administrations of viruses in carrier cells may, therefore, be necessary. Finally, whether or not viral antigens will be presented by carrier cells, and the impact on virus delivery and replication, needs to be explored.

Targeting the tumor microenvironment enhances viral spread and efficacy

The efficacy of virotherapy can be limiting when replication-mediated oncolysis is the sole MOA. Indeed, the tumor microenvironment plays an important role in restricting viral spread and promoting tumor growth. To address this issue, several approaches have been taken. The first is to engineer viral vectors with therapeutic transgenes that target the key components of the tumor microenvironment (for example, the tumor vasculature or matrix). Oncolytic adenovirus encoding relaxin, a matrix-degrading protein, was able to enhance viral spread without causing significant toxicity.¹⁹ Oncolytic HSV encoding dominant-negative fibroblast growth factor receptor or antiangiogenic protein platelet factor-4 led to significant reduction in tumor vasculature and as a result, significantly enhanced therapeutic efficacy.^{20,21} Others have engineered oncolytic virus replication to be activated by tumor matrix metalloproteinases (MMP),²² and shown that MMP-8 gene delivery enhanced the efficacy of oncolytic adenovirus.²³

An alternative approach is to coadminister therapeutic agents with the virus. Coadministration of matrix-modifying agents (bacterial collagenase, MMP-1, 8) has been shown to enhance the spread of oncolytic HSV,^{24,25} although concerns about tumor metastases have to be explored in more preclinical models before translation into clinical trials. In addition, infection with wild-type HSV results in reduction in thrombospondin secretion, a protein that has antiangiogenic properties, from extracellular matrix. This leads to increased vascularity in infected tissues. Aghi *et al.*²⁶ showed that increased vascularity can be counteracted by introduction of certain mutations into oncolytic HSV, or coadministering a thrombospondin-derived peptide 3TSR. A recent study by Kolodkin-Gal *et al.*²⁷ showed that the difference in amount of extracellular matrix between normal colon and colon cancer tissues determined the infectivity and subsequent cytotoxicity of HSV. This phenomenon needs to be further investigated.

Tumor hypoxia and its impact on viral replication have also been studied. Previous reports have shown that hypoxia limits group C adenovirus (serotype 5) replication, and new data suggest that the oncolytic activity of other adenovirus serotypes are also affected.²⁸ Importantly, the report showed that the expression level of CD46, a receptor for group B adenoviruses (serotypes 3, 11) as well as measles virus, was not altered in hypoxic conditions. In contrast, we have found that hypoxia

enhances the replication of oncolytic HSV (M Aghi *et al.*, unpublished).

Another important issue is to explore how inflammation induced by virus infection impacts on the tumor microenvironment. Breitbach *et al.*²⁹ showed that administration with VSV and vaccinia viruses resulted in a dramatic transcriptional activation of the pro-inflammatory neutrophil chemoattractants CXCL1 and CXCL5 and neutrophil attraction. The neutrophils in turn contributed to acute reduction in tumor vasculature. Targeted recruitment of neutrophils to infected tumor beds enhances the killing of malignant cells.²⁹ Recent work by Kurozumi *et al.*³⁰ also illustrates the importance of targeting the tumor microenvironment to improve the efficacy of oncolytic virotherapy. Using the orthotopic (that is, tumors grown in the organs where it is derived from) immunocompetent rat glioma model, they showed that oncolytic HSV infection increased tumor vascular permeability, host leukocyte infiltration into tumors and intratumoral expression of inflammatory cytokine genes, all of these were part of the inflammatory response after HSV infection. Pretreatment with cyclophosphamide suppressed the inflammation and resulted in reduced tumor vascular permeability.³⁰

Kirn *et al.*³¹ showed that systemically administered vaccinia virus resulted in infection and subsequent destruction of tumor endothelial cells, which led to loss of tumor vascular density.

Oncolytic viruses kill cancer stem cells

In light of recent discoveries in the field of cancer stem cells, it is becoming clear that those cell populations not only initiate tumorigenesis, but also contribute importantly to resistance to chemo- and radiation therapy. As this cell population is capable of replication and self-renewal, oncolytic viruses that are designed to target cell cycle-dysregulated tumor cells might also possess the ability to kill cancer stem cells. Indeed, several recent publications have shown that the adenovirus E1A mutant that targets the retinoblastoma-E2F transcriptional factor pathway (Delta-24) is able to kill CD133+ cancer stem cells or CD44(+)/CD22(-/low) cancer initiating cells *in vitro*, and is also able to eradicate tumors derived from these cancer stem cells.^{32,33} MOA include replication-induced cell lysis (necrosis) and autophagy (degradation of intracellular components in lysosomes).³² It has also been reported that adenovirus serotype 3 was better than serotype 5 in infecting cancer stem cells *in vitro*. This ability to kill cancer stem cells does not seem to be limited to adenoviruses, as oncolytic HSV can also efficiently kill glioma stem cells (H Wakimoto *et al.*, unpublished). While this is of interest, there are several issues that remain to be solved. As the population of cancer stem cells within a tumor is generally low (often less than 5%), it is a challenge for oncolytic viruses to 'find' and kill these cancer stem cells, especially when viruses are administered systemically. Secondly, there is so far no direct evidence of anticancer stem cells efficacy of this approach *in vivo* in tumors.

Genetic engineering of oncolytic viruses complements chemo- and molecular-targeted therapies

Several novel combination treatments have been tested in combination with oncolytic viruses. Genetic engineering

of the viruses allows functional complementation to chemotherapeutic agents and molecular-targeted therapeutics. Aghi *et al.*³⁴ showed that temozolomide-induced DNA repair pathways in glioma cells complemented replication of γ 34.5-deleted oncolytic HSV replication and resulted in enhanced efficacy both *in vitro* and *in vivo*. Stanford *et al.*³⁵ and Lun *et al.*³⁶ showed that treatment of rapamycin, a mammalian target of rapamycin inhibitor which resulted in increased Akt/protein kinase B activation, one was able to enhance myxoma virus replication in tumor, but not normal, cells. We showed that by deleting Us3, oncolytic HSV can synergize with Phosphoinositide-3 kinase (PI3K)/Akt inhibitors *in vitro* and have enhanced efficacy without increasing toxicity *in vivo*.³⁷

Histone deacetylase (HDAC) inhibitors are currently being investigated in combination with various oncolytic virus species. Trichostatin A (TSA) has been shown to upregulate cellular coxsackievirus-adenovirus receptor (CAR) expression and hence, infectibility of tumor cells to adenoviruses. We showed that in addition to CAR upregulation, TSA possesses antitumoral and antiangiogenic activities, and shows synergistic tumor-killing effect with oncolytic HSVs *in vitro* and enhanced efficacy *in vivo* (Liu *et al.*³⁸). In contrast, valproic acid, another HDAC inhibitor used for epilepsy disorders, showed antagonistic effect with oncolytic adenoviruses *in vitro*, presumably due to enhanced apoptosis that limits viral replication.³⁹ It will be interesting to see what the effect other types of HDAC inhibitors will have on different oncolytic virus species.

Genetic engineering of oncolytic viruses targets cancer signaling pathways

Through genetic engineering, viruses can be designed to target cancer cells through certain activated signaling pathways. Apart from the Akt pathway targeted by myxoma virus described above, HSV γ 34.5-deleted mutants showed enhanced replication in cells with activated mitogen-activated protein (MAP) kinase or extracellular signal-regulated (ERK) kinase, which in turn inhibited protein kinase R activity, thus circumventing the negative impact of the IFN signaling pathway.^{40,41} Similarly, viruses such as VSV showed preferential replication in cells with an activated Ras-ERK pathway and defective IFN pathways.⁴² A vaccinia virus mutant with a deletion in B18R, whose gene product neutralizes type I IFNs, showed IFN-dependent cancer selectivity and efficacy.³¹ It has also been shown that adenovirus-induced ERK activation is critical to viral replication.⁴³ Oncolytic viruses can also be 'programmed' to replicate in cells through certain cellular signaling activities, such as β -catenin,⁴⁴ to carry therapeutic transgene that targets tumorigenic pathways,²⁰ or retargeted to cellular receptors that are essential for signaling (for example, epidermal growth factor receptor (EGFR)).⁴⁵

Novel oncolytic virus species are being explored

As most oncolytic viruses have produced less than optimal efficacy in clinical trials as single agents, there is great interest in exploring novel viral species. These studies assess oncolytic activity and/or investigate tumor selectivity. For example, the porcine Seneca Valley virus has recently been discovered to possess anti-

tumoral activity against certain cancers of neuroendocrine origin.⁴⁶ It has been speculated that the tumor selectivity is based on differential receptor binding in cancer and normal cells, but more work needs to be done to verify this and to study the impact of the immune system has on the virus. Myxoma virus, a rabbit virus, has also been assessed as an oncolytic agent. New studies reveal that the tumor selectivity of myxoma virus is based on overexpression of Akt in human cancer cells, which facilitates replication and oncolysis.⁴⁷ While these are important findings, the safety profiles of these viruses need to be cautiously examined, especially when the natural host of the virus is not human.

Intratumoral administration of UV-inactivated, replication-deficient Sendai virus induced a robust antitumoral immune response (including cytotoxic T lymphocyte (CTL) induction, dendritic cell maturation and antagonism of Tregs) and resulted in significant efficacy in CT26 syngeneic murine colorectal cancer model.⁴⁸ It will be interesting to study how much of this vaccine effect contributes to antitumoral efficacy in oncolytic Sendai virus studies. A 'vaccine' effect was also seen with parvovirus H-1 in a rat lung tumor metastases model.⁴⁹

A large number of clinical trials have been carried out Virotherapy has several features that are distinct from other therapeutics. Its multiple novel MOAs include replication-mediated oncolysis, antitumoral immunity induction, antiangiogenesis, apoptosis and autophagy induction. There is no cross resistance with other therapeutics, and synergistic interaction is seen with other treatment modalities. Safety in human has been demonstrated in more than 800 patients. In addition, current biotechnology allows us to rapidly address issues encountered in clinics at the bench. There are several reports on virotherapy clinical trials over the past 2 years. Readers are referred to other articles for a more comprehensive review.¹ A list of oncolytic virus agents that have completed, or are currently in late phase trials, are listed in Table 2. In a recent study examining the OV 001 (HUJ) strain of Newcastle disease virus (NDV), OV 001 was administered intravenously to 11 patients with glioblastoma with no dose-limiting toxicity.⁵⁰ Following

biweekly maintenance therapy, one complete remission with a duration of 3 months was described. Virus was recovered from blood, urine and saliva samples. Infectious NDV was recovered from a tumor biopsy. In a separate study testing intravenous delivery of a different NDV strain PV701, using 'two-step' desensitization (dosing with significantly smaller doses prior to full dose) has proven to significantly reduce acute adverse events.⁵¹

The HSV-1 mutant NV1020 (R7020) virus was originally developed as a vaccine. The virus has a deletion in one of the two copies of the γ -34.5 gene. A phase I trial of NV1020 administered by hepatic arterial infusion (HAI) was performed in HSV-seropositive patients with colorectal liver metastases.⁵² No significant toxicity was noted in patients receiving doses up to 1×10^8 pfu per infusion. No replication data were reported. An ongoing phase I/II trial is evaluating repeat HAI of NV1020 followed by second-line chemotherapy in seropositive patients with colorectal cancer. The oncolytic HSV vector OncoVEX^{GM-CSF} has been generated with deletions in γ -34.5 (to reduce pathogenicity) and ICP47 (to restore MHC I presentation). In addition, OncoVEX^{GM-CSF} has a granulocyte monocyte colony-stimulating factor (GM-CSF) transgene insertion. An early passage clinical isolate was used to generate OncoVEX^{GM-CSF} because it had enhanced potency relative to available laboratory strains. A phase I trial of intratumoral injection of OncoVEX^{GM-CSF} into cutaneous metastases from solid tumors and melanomas was carried out.⁵³ Treatment did not result in significant systemic toxicities; injection-site inflammation was dose limiting. Viral genomes were detected shedding from the skin over ulcerated tumors. Injected tumor histology showed inflammation and necrosis. No distant efficacy was reported. Phase II trials of intratumoral injection using OncoVEX^{GM-CSF} are underway in patients with melanoma and other tumor types. A Phase III trial of HSV-1 mutant 1716 in brain tumor has also been announced.

In addition, an oncolytic vaccinia virus with deletion in thymidine kinase (TK) and expressing GM-CSF, JX-594, has been tested in patients with liver tumors. Unlike other virus species described above, systemic exposure and evidence of replication *in vivo* has been

Table 2 Oncolytic viral agents in completed or ongoing late phase trials

Product	Species	Genetic modification	Target tumor type	Phase
Reolysin	Reovirus	None	Bone/soft-tissue sarcoma	II
NDV (MTH-68H)	Newcastle disease virus	None	Metastatic solid tumors	II
JX-594	Vaccinia	Thymidine kinase deletion; GM-CSF expression	1. Hepatocellular 2. Melanoma	II
H101	Adenovirus	E1B-55 K, E3 deletion	Head and neck (+ chemotherapy)	III
Ad5-yCD/ mutTKSR39rep-ADP	Adenovirus	E1B-55 K deletion; CD/TK fusion gene expression; ADP overexpression	Prostate (+ radiotherapy)	II
OncoVex ^{GM-CSF}	HSV 1	γ 34.5 and ICP47 deletion; GM-CSF expression	Melanoma	II
1716	HSV 1	γ 34.5 deletion	Brain	III

Abbreviations: ADP, adenovirus death protein; GM-CSF, granulocyte monocyte colony-stimulating factor; HSV, herpes simplex virus; TK, thymidine kinase; VSV, vesicular stomatitis virus.

demonstrated in this study. Tumor response was observed in the majority of treated tumors, and distant tumor responses were demonstrated in several patients with target tumor responses. Replication and biological activity *in vivo* have also been shown.⁵⁴ With the advance in our knowledge of cancer and oncolytic virus biology, we expect that carefully designed clinical studies will not only show proof-of-concept for this novel treatment, but also result in evidence of clinical benefit in patients.

Prospects

Logical design of the next generation of oncolytic viruses may take this strategy to the next level

Design of the next generation of oncolytic viruses should be based on current knowledge of virology, immunology and cancer biology. More importantly, findings from clinical trials should be incorporated/addressed. Two recent publications illustrate this concept.^{31,55} The first article describes rational species and strain selection and genetic engineering based on updated knowledge. As described before, poxvirus was selected for this study based on human data showing that systemic delivery of poxvirus is safe and can induce significant tumor responses.⁵⁶ A highly potent vaccinia virus strain that also trafficked efficiently to human tumors after intravenous administration was first identified. This strain was then engineered to target cancer cells with activated transcription factor E2F and the EGFR pathway, and further engineered to express human GM-CSF for induction of tumor-specific CTL. The new vaccinia construct, JX-963, demonstrated significant cancer selectivity in human tumor cell lines, tumor-bearing rabbits and primary human surgical samples. Intravenous administration led to systemic efficacy against both primary carcinomas and widespread organ-based metastases in immunocompetent animals.⁵⁵ The second study describes the use of a vaccinia virus background that selectively targets IFN pathway resistance in tumor cells. Further engineering with TK deletion and IFN- β insertion results in a multimechanistic oncolytic vaccinia virus.³¹

Ex vivo studies may predict responses

Genetic markers are being developed for chemotherapy and molecular-targeted therapeutics to predict responses and/or idiosyncratic reactions, and the results have been implemented into practice. Given the complexity of MOA of virotherapeutics, there is therefore a long way to go before such markers/predictors can be developed for virotherapy. However, testing patients' samples *ex vivo* prior to treatment may provide important information and complement our current studies. In addition to the difference between results obtained from immortalized cell lines and primary cancer cells, explant samples contain extracellular matrix and a three-dimensional structure that more closely mimics clinical situation. Importantly, when adjacent normal tissues are included, a therapeutic index can be obtained, which is critical for local administration protocol. Several recent publications described the use of this approach.^{27,55,57,58} The results of oncolytic viruses on *ex vivo* tissues and its clinical outcome correlation have yet to be established, but researchers are encouraged to include explants

studies whenever possible to maximize clinical relevance. Future clinical developments might include pre-/post-treatment *ex vivo* assessment of infectivity, cytopathic effects and viral replication.

The demand for target validation is increasing

The demand for target validation in small molecule-based kinase inhibitors is increasing. For virotherapy to be successful, it has to pass similar hurdles. The most important biological endpoint that needs to be demonstrated with all species of oncolytic virus is tumor-selective virus replication, therapeutic transgene expression and biological function (if applicable). This has been effectively achieved with several virus species, but still required for others. Depending on the strategy demonstration of targeting cancer-specific features pathways may also be necessary. For instance, if viruses are designed to target the Ras/Raf/MAPK pathway, a reduction in pathway activity should be proven. High TK activity in cancer cells are needed for TK-deleted viruses, while viruses replicating exclusively in IFN-resistant cells (cancer cells) need to demonstrate local IFN induction *in vivo*. These are difficult tasks but warrant further exploration. The effect of oncolytic viruses on tumor microenvironment, as shown in various preclinical studies, will need to be validated in clinical trials. For instance, future clinical studies should include tumor vascularity assessment to see if virotherapy reduces tumor vasculature.

New imaging endpoints in clinical trials?

Tumor size measurement has been a gold standard for defining responses in clinical practice. Tumor progression is defined as increased in tumor size above certain degrees. However, recent studies on molecular therapeutics have shown that some agents induce tumor necrosis without affecting the sizes of the tumors. To address this issue, new criteria (for example, Choi criteria⁵⁹) have been incorporated in tumor response assessment by imaging. Since most of the virotherapy clinical trials were done by local or locoregional administration, it is likely that the effect of the viruses is localized within the tumors. Thus, we need to consider whether adopting these new criteria is feasible. Correlation between tumor size, tumor density and survival will be necessary in late phase trials.

Abbreviations

CAR, coxsackievirus-adenovirus receptor; CIK, cytokine-induced killer; CTL, cytotoxic T lymphocyte; EGFR, epidermal growth factor receptor; HDAC, histone deacetylase; HSV, herpes simplex virus; IFN, interferon; MMP, matrix metalloproteinase; MOA, mechanism-of-action; PBMC, peripheral blood mononuclear cells; TK, thymidine kinase; Treg, regulatory T cells; TSA, trichostatin A; VSV, vesicular stomatitis virus

References

- 1 Liu TC, Galanis E, Kirn D. Clinical trial results with oncolytic virotherapy: a century of promise, a decade of progress. *Nat Clin Pract Oncol* 2007; 4: 101–117.

- 2 Fulci G, Breyman L, Gianni D, Kurozumi K, Rhee SS, Yu J *et al.* Cyclophosphamide enhances glioma virotherapy by inhibiting innate immune responses. *Proc Natl Acad Sci USA* 2006; **103**: 12873–12878.
- 3 Qiao J, Wang H, Kottke T, White C, Twigger K, Diaz RM *et al.* Cyclophosphamide facilitates antitumor efficacy against subcutaneous tumors following intravenous delivery of reovirus. *Clin Cancer Res* 2008; **14**: 259–269.
- 4 Lamfers ML, Fulci G, Gianni D, Tang Y, Kurozumi K, Kaur B *et al.* Cyclophosphamide increases transgene expression mediated by an oncolytic adenovirus in glioma-bearing mice monitored by bioluminescence imaging. *Mol Ther* 2006; **14**: 779–788.
- 5 Fulci G, Dmitrieva N, Gianni D, Fontana EJ, Pan X, Lu Y *et al.* Depletion of peripheral macrophages and brain microglia increases brain tumor titers of oncolytic viruses. *Cancer Res* 2007; **67**: 9398–9406.
- 6 Li H, Zeng Z, Fu X, Zhang X. Coadministration of a herpes simplex virus-2 based oncolytic virus and cyclophosphamide produces a synergistic antitumor effect and enhances tumor-specific immune responses. *Cancer Res* 2007; **67**: 7850–7855.
- 7 Ungerechts G, Springfield C, Frenzke ME, Lampe J, Parker WB, Sorscher EJ *et al.* An immunocompetent murine model for oncolysis with an armed and targeted measles virus. *Mol Ther* 2007; **15**: 1991–1997.
- 8 Di Paolo NC, Tuve S, Ni S, Hellstrom KE, Hellstrom I, Lieber A. Effect of adenovirus-mediated heat shock protein expression and oncolysis in combination with low-dose cyclophosphamide treatment on antitumor immune responses. *Cancer Res* 2006; **66**: 960–969.
- 9 Haralambieva I, Iankov I, Hasegawa K, Harvey M, Russell SJ, Peng KW. Engineering oncolytic measles virus to circumvent the intracellular innate immune response. *Mol Ther* 2007; **15**: 588–597.
- 10 Zhu H, Su Y, Zhou S, Xiao W, Ling W, Hu B *et al.* Immune analysis on mTHSV mediated tumor therapy in HSV-1 seropositive mice. *Cancer Biol Ther* 2007; **6**: 724–731.
- 11 Diaz RM, Galivo F, Kottke T, Wongthida P, Qiao J, Thompson J *et al.* Oncolytic immunovirotherapy for melanoma using vesicular stomatitis virus. *Cancer Res* 2007; **67**: 2840–2848.
- 12 Thorne SH, Negrin RS, Contag CH. Synergistic antitumor effects of immune cell-viral biotherapy. *Science* 2006; **311**: 1780–1784.
- 13 Power AT, Bell JC. Cell-based delivery of oncolytic viruses: a new strategic alliance for a biological strike against cancer. *Mol Ther* 2007; **15**: 660–665.
- 14 Ong HT, Hasegawa K, Dietz AB, Russell SJ, Peng KW. Evaluation of T cells as carriers for systemic measles virotherapy in the presence of antiviral antibodies. *Gene Therapy* 2007; **14**: 324–333.
- 15 Power AT, Wang J, Falls TJ, Paterson JM, Parato KA, Lichty BD *et al.* Carrier cell-based delivery of an oncolytic virus circumvents antiviral immunity. *Mol Ther* 2007; **15**: 123–130.
- 16 Iankov ID, Blechacz B, Liu C, Schmeckpeper JD, Tarara JE, Federspiel MJ *et al.* Infected cell carriers: a new strategy for systemic delivery of oncolytic measles viruses in cancer virotherapy. *Mol Ther* 2007; **15**: 114–122.
- 17 Hakkarainen T, Sarkioja M, Lehenkari P, Miettinen S, Ylikomi T, Suuronen R *et al.* Human mesenchymal stem cells lack tumor tropism but enhance the antitumor activity of oncolytic adenoviruses in orthotopic lung and breast tumors. *Hum Gene Ther* 2007; **18**: 627–641.
- 18 Qiao J, Kottke T, Willmon C, Galivo F, Wongthida P, Diaz RM *et al.* Purging metastases in lymphoid organs using a combination of antigen-nonspecific adoptive T cell therapy, oncolytic virotherapy and immunotherapy. *Nat Med* 2008; **14**: 37–44.
- 19 Kim JH, Lee YS, Kim H, Huang JH, Yoon AR, Yun CO. Relaxin expression from tumor-targeting adenoviruses and its intratumoral spread, apoptosis induction, and efficacy. *J Natl Cancer Inst* 2006; **98**: 1482–1493.
- 20 Liu TC, Zhang T, Fukuhara H, Kuroda T, Todo T, Canron X *et al.* Dominant-negative fibroblast growth factor receptor expression enhances antitumoral potency of oncolytic herpes simplex virus in neural tumors. *Clin Cancer Res* 2006; **12**: 6791–6799.
- 21 Liu TC, Zhang T, Fukuhara H, Kuroda T, Todo T, Martuza RL *et al.* Oncolytic HSV armed with platelet factor 4, an antiangiogenic agent, shows enhanced efficacy. *Mol Ther* 2006; **14**: 789–797.
- 22 Springfield C, von Messling V, Frenzke M, Ungerechts G, Buchholz CJ, Cattaneo R. Oncolytic efficacy and enhanced safety of measles virus activated by tumor-secreted matrix metalloproteinases. *Cancer Res* 2006; **66**: 7694–7700.
- 23 Cheng J, Sauthoff H, Huang Y, Kutler DI, Bajwa S, Rom WN *et al.* Human matrix metalloproteinase-8 gene delivery increases the oncolytic activity of a replicating adenovirus. *Mol Ther* 2007; **15**: 1982–1990.
- 24 McKee TD, Grandi P, Mok W, Alexandrakis G, Insin N, Zimmer JP *et al.* Degradation of fibrillar collagen in a human melanoma xenograft improves the efficacy of an oncolytic herpes simplex virus vector. *Cancer Res* 2006; **66**: 2509–2513.
- 25 Mok W, Boucher Y, Jain RK. Matrix metalloproteinases-1 and -8 improve the distribution and efficacy of an oncolytic virus. *Cancer Res* 2007; **67**: 10664–10668.
- 26 Aghi M, Rabkin SD, Martuza RL. Angiogenic response caused by oncolytic herpes simplex virus-induced reduced thrombospondin expression can be prevented by specific viral mutations or by administering a thrombospondin-derived peptide. *Cancer Res* 2007; **67**: 440–444.
- 27 Kolodkin-Gal D, Zamir G, Pikarski E, Pikarski A, Shimony N, Wu H *et al.* A novel system to study adenovirus tropism to normal and malignant colon tissues. *Virology* 2007; **357**: 91–101.
- 28 Shen BH, Bauzon M, Hermiston TW. The effect of hypoxia on the uptake, replication and lytic potential of group B adenovirus type 3 (Ad3) and type 11p (Ad11p). *Gene Therapy* 2006; **13**: 986–990.
- 29 Breitbach CJ, Paterson JM, Lemay CG, Falls TJ, McGuire A, Parato KA *et al.* Targeted inflammation during oncolytic virus therapy severely compromises tumor blood flow. *Mol Ther* 2007; **15**: 1686–1693.
- 30 Kurozumi K, Hardcastle J, Thakur R, Yang M, Christoforidis G, Fulci G *et al.* Effect of tumor microenvironment modulation on the efficacy of oncolytic virus therapy. *J Natl Cancer Inst* 2007; **99**: 1768–1781.
- 31 Kim DH, Wang Y, Le Boeuf F, Bell J, Thorne SH. Targeting of interferon-beta to produce a specific, multi-mechanistic oncolytic vaccinia virus. *PLoS Med* 2007; **4**: e353.
- 32 Jiang H, Gomez-Manzano C, Aoki H, Alonso MM, Kondo S, McCormick F *et al.* Examination of the therapeutic potential of Delta-24-RGD in brain tumor stem cells: role of autophagic cell death. *J Natl Cancer Inst* 2007; **99**: 1410–1414.
- 33 Eriksson M, Guse K, Bauerschmitz G, Virkkunen P, Tarkkanen M, Tanner M *et al.* Oncolytic adenoviruses kill breast cancer initiating CD44(+)CD24(-/low) cells. *Mol Ther* 2007; **15**: 2088–2093.
- 34 Aghi M, Rabkin S, Martuza RL. Effect of chemotherapy-induced DNA repair on oncolytic herpes simplex viral replication. *J Natl Cancer Inst* 2006; **98**: 38–50.
- 35 Stanford MM, Barrett JW, Nazarian SH, Werden S, McFadden G. Oncolytic virotherapy synergism with signaling inhibitors: rapamycin increases myxoma virus tropism for human tumor cells. *J Virol* 2007; **81**: 1251–1260.
- 36 Lun XQ, Zhou H, Alain T, Sun B, Wang L, Barrett JW *et al.* Targeting human medulloblastoma: oncolytic virotherapy with myxoma virus is enhanced by rapamycin. *Cancer Res* 2007; **67**: 8818–8827.
- 37 Liu TC, Wakimoto H, Martuza RL, Rabkin SD. Herpes simplex virus us3(-) mutant as oncolytic strategy and synergizes with phosphatidylinositol 3-kinase-akt targeting molecular therapeutics. *Clin Cancer Res* 2007; **13**: 5897–5902.

- 38 Liu TC, Castelo-Branco P, Rabkin SD, Martuza RL. Trichostatin A and oncolytic HSV combination therapy shows enhanced antitumoral and antiangiogenic effects. *Mol Ther* 2008; e-pub ahead of print 1 April 2008; doi:10.1038/mt.2008.33.
- 39 Hoti N, Chowdhury W, Hsieh JT, Sachs MD, Lupold SE, Rodriguez R. Valproic acid, a histone deacetylase inhibitor, is an antagonist for oncolytic adenoviral gene therapy. *Mol Ther* 2006; **14**: 768–778.
- 40 Smith KD, Mezhr JJ, Bickenbach K, Veerapong J, Charron J, Posner MC *et al.* Activated MEK suppresses activation of PKR and enables efficient replication and *in vivo* oncolysis by Deltagamma(1)34.5 mutants of herpes simplex virus 1. *J Virol* 2006; **80**: 1110–1120.
- 41 Veerapong J, Bickenbach KA, Shao MY, Smith KD, Posner MC, Roizman B *et al.* Systemic delivery of (gamma1)34.5-deleted herpes simplex virus-1 selectively targets and treats distant human xenograft tumors that express high MEK activity. *Cancer Res* 2007; **67**: 8301–8306.
- 42 Noser JA, Mael AA, Sakuma R, Ohmine S, Marcato P, Lee PW *et al.* The RAS/Raf1/MEK/ERK signaling pathway facilitates VSV-mediated oncolysis: implication for the defective interferon response in cancer cells. *Mol Ther* 2007; **15**: 1531–1536.
- 43 Schumann M, Dobbelsstein M. Adenovirus-induced extracellular signal-regulated kinase phosphorylation during the late phase of infection enhances viral protein levels and virus progeny. *Cancer Res* 2006; **66**: 1282–1288.
- 44 Kuroda T, Rabkin SD, Martuza RL. Effective treatment of tumors with strong beta-catenin/T-cell factor activity by transcriptionally targeted oncolytic herpes simplex virus vector. *Cancer Res* 2006; **66**: 10127–10135.
- 45 Allen C, Vongpunsawad S, Nakamura T, James CD, Schroeder M, Cattaneo R *et al.* Retargeted oncolytic measles strains entering via the EGFRvIII receptor maintain significant antitumor activity against gliomas with increased tumor specificity. *Cancer Res* 2006; **66**: 11840–11850.
- 46 Reddy PS, Burroughs KD, Hales LM, Ganesh S, Jones BH, Idamakanti N *et al.* Seneca Valley virus, a systemically deliverable oncolytic picornavirus, and the treatment of neuroendocrine cancers. *J Natl Cancer Inst* 2007; **99**: 1623–1633.
- 47 Wang G, Barrett JW, Stanford M, Werden SJ, Johnston JB, Gao X *et al.* Infection of human cancer cells with myxoma virus requires Akt activation via interaction with a viral ankyrin-repeat host range factor. *Proc Natl Acad Sci USA* 2006; **103**: 4640–4645.
- 48 Kurooka M, Kaneda Y. Inactivated Sendai virus particles eradicate tumors by inducing immune responses through blocking regulatory T cells. *Cancer Res* 2007; **67**: 227–236.
- 49 Raykov Z, Grekova S, Galabov AS, Balboni G, Koch U, Aprahamian M *et al.* Combined oncolytic and vaccination activities of parvovirus H-1 in a metastatic tumor model. *Oncol Rep* 2007; **17**: 1493–1499.
- 50 Freeman AI, Zakay-Rones Z, Gomori JM, Linetsky E, Rasooly L, Greenbaum E *et al.* Phase I/II trial of intravenous NDV-HUJ oncolytic virus in recurrent glioblastoma multiforme. *Mol Ther* 2006; **13**: 221–228.
- 51 Laurie SA, Bell JC, Atkins HL, Roach J, Bamat MK, O'Neil JD *et al.* A phase 1 clinical study of intravenous administration of PV701, an oncolytic virus, using two-step desensitization. *Clin Cancer Res* 2006; **12**: 2555–2562.
- 52 Kemeny N, Brown K, Covey A, Kim T, Bhargava A, Brody L *et al.* Phase I, open-label, dose-escalating study of a genetically engineered herpes simplex virus, NV1020, in subjects with metastatic colorectal carcinoma to the liver. *Hum Gene Ther* 2006; **17**: 1214–1224.
- 53 Hu JC, Coffin RS, Davis CJ, Graham NJ, Groves N, Guest PJ *et al.* A phase I study of OncoVEXGM-CSF, a second-generation oncolytic herpes simplex virus expressing granulocyte macrophage colony-stimulating factor. *Clin Cancer Res* 2006; **12**: 6737–6747.
- 54 Park BH, Hwang TH, Kim SG, Rhee BG, Ahn YJ, Kwon HC *et al.* A phase I-II clinical trial with JX-594, a targeted and GM-CSF-armed oncolytic poxvirus, by intratumoral injection in patients with liver tumors. *AACR-NCI-EORTC International Conference: Molecular Targets and Cancer Therapeutics*. American Association for Cancer Research: San Francisco, CA, 2007, pp 116.
- 55 Thorne SH, Hwang TH, O'Gorman WE, Bartlett DL, Sei S, Kanji F *et al.* Rational strain selection and engineering creates a broad-spectrum, systemically effective oncolytic poxvirus, JX-963. *J Clin Invest* 2007; **117**: 3350–3358.
- 56 Liu TC, Kirn D. Systemic efficacy with oncolytic virus therapeutics: clinical proof-of-concept and future directions. *Cancer Res* 2007; **67**: 429–432.
- 57 Rots MG, Elferink MG, Gommans WM, Oosterhuis D, Schalk JA, Curiel DT *et al.* An *ex vivo* human model system to evaluate specificity of replicating and non-replicating gene therapy agents. *J Gene Med* 2006; **8**: 35–41.
- 58 Wollmann G, Robek MD, van den Pol AN. Variable deficiencies in the interferon response enhance susceptibility to vesicular stomatitis virus oncolytic actions in glioblastoma cells but not in normal human glial cells. *J Virol* 2007; **81**: 1479–1491.
- 59 Choi H, Charnsangavej C, Faria SC, Macapinlac HA, Burgess MA, Patel SR *et al.* Correlation of computed tomography and positron emission tomography in patients with metastatic gastrointestinal stromal tumor treated at a single institution with imatinib mesylate: proposal of new computed tomography response criteria. *J Clin Oncol* 2007; **25**: 1753–1759.